

Quantum chemical analysis of the hydrogen bond interactions in secondary structures of proteins with Negative Fragmentation Approach

Yu Takano^{1,2}, Hiroko X. Kondo^{1,3}, and Haruki Nakamura²

¹ Graduate School of Information Sciences, Hiroshima City University, Hiroshima 731-3194 Japan

² Institute for Protein Research, Osaka University, Suita 565-0871 Japan

³ Faculty of Engineering, Kitami Institute of Technology, Kitami, 090-8507, Japan

Hydrogen bond (H-bond) is one of the major factors that build the macromolecular structures of proteins, nucleic acids and their complexes. In particular, pair-wise H-bonds in protein backbones are essential to form their characteristic three-dimensional (3D) structures based on their ordered secondary structures, α -helices and β -sheets. Therefore, their structural energies should be correctly computed for analyses and predictions of protein 3D structures.

In this study, we have investigated the hydrogen bond interactions in secondary structures, α -helices, β -sheets, and turns by precise DFT calculations, followed by a negative fragmentation approach to extract a local interaction between molecular fragments [1–3]. Our computation showed that the reduced H-bond energy in α -helix originated from the depolarizations of both the H-bond donor and acceptor groups, due to the repulsive interactions with the neighboring polar peptide groups in the α -helix backbone [1].

References

- [1] H. X. Kondo, A. Kusaka, C. K. Kitakawa, J. Onari, S. Yamanaka, H. Nakamura, Y. Takano, *J. Comput. Chem.* **2019**, *40*, 2043–2052.
- [2] Y. Takano, A. Kusaka, H. Nakamura, *Biophys. Physicobiol.* **2016**, *13*, 27–35.
- [3] M. M. Deshmukh, S. R. Gadre, *J. Phys. Chem. A* **2009**, *113*, 7927–7932.