

Accurate van-der-Waals interactions from (semi)-local density functional theory

Alexandre Tkatchenko* and Matthias Scheffler

Fritz-Haber-Institut der Max-Planck-Gesellschaft, Faradayweg 4-6, 14195, Berlin, Germany

Non-covalent forces, such as hydrogen bonding and van der Waals (vdW) interactions, are crucial for the formation, stability and function of molecules and materials. At present, vdW interactions can only be satisfactorily accounted for by high-level quantum-chemical wave function or by the Quantum Monte Carlo (QMC) method. In contrast, (semi)-local DFT and Hartree-Fock approximation fail for the description of vdW forces. We present a parameter-free method for describing the long-range vdW interaction in (semi)-local DFT. The leading C_6 coefficients are derived from the electron density of a molecule/solid and accurate reference values for the free atoms. The mean absolute error in the C_6 coefficients is 5.5% when compared to accurate experimental values for 1225 intermolecular pairs. Furthermore, the results are essentially independent of the employed (semi)-local exchange-correlation functional. We show that the C_6 coefficients depend strongly on the bonding type and geometry of molecules/solids. Finally, we analyze the vdW radii and the damping function in the C_6R^{-6} correction method for DFT calculations.

* Electronic address: tkatchen@fhi-berlin.mpg.de