

CAS-DFT based on Coulomb-division scheme

S. Yamanaka^a, T. Ukai^b, K. Nakata^b, K. Kusakabe^c,
Y. Yonezawa^d, H. Nakamura^d, T. Takada^e, and K. Yamaguchi^a

^a Center for Quantum Science and Technology under Extreme Conditions, Osaka University, Toyonaka, 560-8531, Japan.

^b Department of Chemistry, Graduate School of Science, Osaka University, Toyonaka, 560-0043, Japan.

^c Department of Physical Science, Graduate School of Engineering Science, Osaka University, Toyonaka, 560-8531, Japan

^d Institute for Protein Research, Osaka University, Toyonaka, 560-8531, Japan

^e RIKEN, Next-Generation Supercomputer R&D center, Chiyodaku, Tokyo, 100-0005, Japan

During the past decade, a new class of electronic structure theory combining multireference wavefunction and density functional (MR-DFT) has been developed to cover the systems that exhibit strong multireference character [1-5]. There are various types of MR-DFT, which are classified into Coulomb-driven [1-3] and wavefunction-driven ones [3-5]. The former is based on the division of electron repulsion Coulomb operator into short-and-long range parts. On the other hand, wavefunction-driven MR-DFT is based on truncation of wavefunction expansions, and remaining correlations are covered by DFT. From the computational viewpoint, a modest wavefunction (WF), such as complete-active-space (CAS) WF, is desirable. Indeed, we present CASCI/CASSCF-DFT methods that satisfy the variational principle for both ground and excited states [3,5]. In these previous studies, we employ MSS-GC scheme [4] for CAS-DFT. In this study, we propose a new type of MR-DFT, in which a long or short-range part of the Coulomb repulsions for nondynamical correlations is covered by CAS wavefunction and other all parts are covered by DFT. Numerical examples are presented for both ground-and-excited states of simple systems.

[1] A. Savin, H-J. Flad, *Int. J. Quantum Chem.*, **56**, 327(1995); P. Gori-Giorgi, A. Savin, *Phys. Rev. A* **73**(2006)3250; E. Fromager, et al. *J. Chem. Phys.* **126** 74111(2007).

[2] K. Kusakabe, *J. Phys. Soc. Jpn* **70**, 2038(2001).

[3] S. Yamanaka et al. *Chem. Lett.* **35**, 242(2006); *Int J Quantum Chem.* **16**, 3312(2006).

[4] B. Miehlich, H. Stoll, A. Savin, *Mol. Phys.*, **91**, 527 (1997); J. Gräfenstein, D. Cremer, *Chem. Phys. Lett.* **316**, 569 (2000); S. Gusarov et al. *Theor. Chem. Acc.* **112**, 84(2004); A. J. Pérez-Jiménez, J. M. Pérez-Jord, *Phys. Rev.* **A75**, 012503 (2007); R. Takeda, et al., *Chem. Phys. Lett.* **366**, 321(2002); *Int. J Quantum Chem.* **96**, 463 (2004).

[5] T. Ukai et al. *Mol. Phys.* in press; K. Nakata, et al. *AIP Conf. Proc.*, in press.