

Current-density-functional theory of molecules in weak and strong magnetic fields

Trygve Helgaker*, Alex Borgoo*, Ulf E. Ekström*, James Furness†, Jürgen Gauss**, Simen Kvaal*, Kai Lange*, Sarah Reimann*, Espen Sagvolden*, Stella Stopkowicz*, Erik Tellgren* and Andy M. Teale*,‡

*Department of Chemistry University of Oslo, NO-0315 Oslo, Norway

†School of Chemistry, University of Nottingham, University Park, Nottingham, NG7 2RD, UK

**Institut für Physikalische Chemie, Universität Mainz, D-55099 Mainz, Germany

‡School of Chemistry, University of Nottingham, University Park, Nottingham, NG7 2RD, UK.

Abstract. We have undertaken a quantum-mechanical study of molecules in finite magnetic fields, using a recently developed computer code London for the calculation of many-electron molecules with wave-function theory [Tellgren *et al.*, *J. Chem. Phys.* **129**, 154114 (2008)] and more recently with Kohn–Sham density-functional theory (DFT) and current-density-functional theory (CDFT). These two approaches to electronic-structure theory are connected by an implementation of the Lieb variation principle [Teale *et al.*, *J. Chem. Phys.* **130**, 104111 (2009); **132**, 164115 (2010)], allowing us to study the exchange-correlation functional of DFT at different levels of ab initio theory.

Our calculations on molecular systems in strong magnetic fields have revealed and highlighted many interesting phenomena such as the transition to diamagnetism of paramagnetic molecules at a critical field strength [Tellgren *et al.*, *Phys. Chem. Chem. Phys.* **11**, 5489 (2009)]. Perhaps most interestingly, antibonding molecular orbitals are stabilised in the field, leading to strongly bound triplet H₂ and singlet He₂ molecules [Lange *et al.*, *Science* **337**, 327 (2012)] and helium clusters [Tellgren *et al.*, *Phys. Chem. Chem. Phys.* **14**, 9492 (2012)] at field strengths of about 10⁵ Tesla.

The application of DFT to magnetic fields requires modifications of the standard theory, by introducing a field- or current-dependence in the exchange-correlation functional. However, essentially all applications of Kohn–Sham theory to study magnetic phenomena ignore these modifications—yielding, for example, NMR shielding constants that are unreliable [Teale *et al.*, *J. Chem. Phys.* **138**, 024111 (2013)]. We discuss here some aspects of DFT in magnetic fields, including the Hohenberg–Kohn theorem and the choice of variables in DFT [Tellgren *et al.*, *Phys. Rev. A* **86**, 062506 (2012)]. Kohn–Sham calculations with different parameterisations of the Vignale–Rasolt–Geldart (VRG) functional in finite magnetic fields are presented. The results demonstrate that this functional form does not provide improvements in practical calculations and highlight the need for the development of new current-dependent approximations. On the other hand, the TPSS meta-generalised-gradient-approximation (meta-GGA) with a dependence on the kinetic energy density provides a natural framework for introducing a current dependence [Dobson, *J. Chem. Phys.* **98**, 8870 (1993)], giving good results in comparison with high-level many-body methods (FCI and CCSD(T)). In the figure below, dissociation curves of Ne₂ in a strong magnetic field have been plotted. The Tao-Perdew-Staroverov-Scuseria (TPSS and TPSSh) curves with current dependence compare well with CCSD(T) curve.

